

- الوراثة البشرية | المرحلة الثالثة العام الدراسي ٢٠٢٠ ٢٠٢١
- المحاضرة السادسة | التغييرات الصبغية الكروموسومية التركيبية ( Chromosomes Abnormalities ) ( Structural )
  - المحاضر \ الأستاذ الدكتور اياد احمد الطويل
  - كلية الأسراء الجامعة | قسم تقنيات المختبرات الطبية
    - وزارة التعليم العالي و البحث العلمي



#### **Deficiencies**


- Phenotypic consequences of deficiency depends on
  - Size of the deletion
  - Functions of the genes deleted
- Phenotypic effect of deletions usually detrimental



Two types of deletion

#### **Deletion**

- o Loss of a region of chromosome
- A chromosomal deficiency occurs when a chromosome breaks and a fragment is lost



- Known disorders in humans include
- Wolf-Hirschhorn syndrome, which is caused by partial deletion of the short arm of chromosome 4; and
- <u>Jacobsen syndrome</u>, also called the terminal 11q deletion disorder.

#### **Cri-du-chat Syndrome**

- High-pitched cry
- intellectual disability
- delayed development
- small head size
- low birth weight
- weak muscle tone in infant



#### Duplication

- A duplication is a chromosomal mutation that results when a segment of DNA breaks off and attaches onto the homologous chromosome.
- There are different forms of chromosomal duplications or different tandem configuration
  - 1. tandem
  - 2. reverse tandem
  - 3. terminal tandem






Fig. 8.7 Chromosome of Drosophila strains





- <u>Translocations</u>: A portion of one chromosome is transferred to another chromosome. There are two main types of translocations:
- <u>Reciprocal translocation</u>: Segments from two different chromosomes have been exchanged.
- Robertsonian translocation: An entire chromosome has attached to another at the centromere - in humans these only occur with chromosomes 13, 14, 15, 21 and 22.

#### **Reciprocal Translocation**

- Involves two chromosomes
- One break in each chromosome
- The two chromosomes exchange broken segments










#### **Robertsonian Translocation**

- Named after W. R. B. Robertson who first identified them in grasshoppers in 1916
- Most common structural chromosome abnormality in humans
  - Frequency = 1/1000 livebirths



- Involves two acrocentric chromosomes
- Two types
  - Homologous acrocentrics involved
  - Non-Homologous acrocentrics involved









#### Inversion

- A segment of chromosome that is flipped relative to that in the homologue
- Two breaks in one chromosome

The fragment generated rotates 180° and reinserts into the




If the centromere is not part of the rearranged chromosome segment, it is a paracentric inversion.




If the centromere is part of the inverted segment, it is described as a pericentric inversion.

### Isochromosome (Transverse centromeric division)



#### Example:

- \*Long arm of X- chromosome remains, short arms lost
- •Cytogenetic variant of Turner's syndrome




#### <u>Isochromosome</u>

(Transverse centromeric division)

38

37

#### Cont....



#### Dicentric chromosomos


Unlike normal chromosomes, which have a single constriction point (centromere), a dicentric chromosome contains two centromeres. Dicentric chromosomes result from the abnormal fusion of two chromosome pieces, each of which includes a centromere. These structures are unstable and often involve a loss of some genetic material.

#### Cont.....

#### Ring chromosomos:

Ring chromosomes usually occur when a chromosome breaks in two places and the ends of the chromosome arms fuse together to form a circular structure. The ring may or may not include the chromosome's constriction point (centromere). In many cases, genetic material near the ends of the


chromosome is lost.



#### Figure 8.13

Copylight © The McGraw-Hill Companies, Inc. Permission regards for reproduction or display.

Translocation cross



Chromosomal translocations play a large role in blood • cancers. It's estimated that translocation are presented in up to 90% of LYMPHOMAS and over 50% of LEUKEMIAS.

#### Following some examples:

- t (8;22): Acute myeloblastic leukemyia
- t (9;22): Philadelphia Chromosome
- t (15;17): Acute promyelotic leukemyia
- t (12;15), t(1;12) Acute myelogenous leukemyia
- t (2;5) Anaplastic large cell lymphoma
- t (8;14) Burkitt's lymphoma
- t (11;14) Mantle cell lymphoma
- t (14;18) Follicular lymphoma

## SOME Diseases resulted from chromosomal deletion

Cri-du-chat (cat cry) Syndrame 46,XX/XY,5p15.2 / 46,XX/XY,5p15.3

Retinoblastoma 46,XX/XY,13q14

Wilms Tumor 46,XX/XY, 11P13

Wolf-Hirschhorn Syndrome 46, XX/XY, 4 short arm

Williams Syndrome 46,XX/XY, 4 long arm

Prader- Willi Syndrome 46,XX/XY,15 long arm (q11-q13) –

**Paternal** 

Angelman Syndeome 46,XX/XY, 15 long arm(q11-q13) --

**Maternal** 

#### **SOME types of chromosomal duplication**

Duplication of chromosome 4, short arm 46,XX/XY

Duplication of chromosome 4, long arm 46, XX/XY

Duplication of chromosome 7, long arm 46, XX/XY

Duplication of chromosome 9, short arm 46, XX/XY

#### Some types of chromosomal inversion/ below are 3

#### types of inversion in human chromosome

Chromosomal inversion in human is known as inv(9)(p12q13) ( Paracentric inversion }

Chromosomal inversion in human is known as inv(3)(q24q27) { Pricentric inversion }

Chromosomal Inversion in human as inv(2)(p11.2q13) { Paracentric inversion }

#### **IMPORTANT NOTE**

- في الملزمة الصفحتين ٢١ و ٣١ توجد ثلاثة جداول تتضمن عدد من التغييرات الصبغية الكروموسومية العددية و التركيبية و تأثيراتها في الانسان ، مثلا الجدول في صفحة ٢١ يوضح التغييرات الصبغية الكروموسومية التركيبية خصوصا النوعين الفقدان و التضاعف و ماهي الاعراض التي تحدث عنها في الانسان.
- اما الجدولين في الصفحة ٣٤ فيوضحان (الجدول الاول يبين المختصرات و ماذا تعني و الجدول الثاني يبين نوع التغيير و مفهومه)

# Thanks for your listening Dr. Ayad